



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07K 7/06, A61K 37/02 G01N 33/564, 33/68	A1	(11) International Publication Number: WO 90/11294 (43) International Publication Date: 4 October 1990 (04.10.90)									
(21) International Application Number: PCT/US90/01516 (22) International Filing Date: 21 March 1990 (21.03.90) (30) Priority data: <table border="0"> <tr> <td>326,314</td> <td>21 March 1989 (21.03.89)</td> <td>US</td> </tr> <tr> <td>382,085</td> <td>18 July 1989 (18.07.89)</td> <td>US</td> </tr> <tr> <td>382,086</td> <td>18 July 1989 (18.07.89)</td> <td>US</td> </tr> </table> (71) Applicant: THE IMMUNE RESPONSE CORPORATION [US/US]; 6455 Nancy Ridge Drive, San Diego, CA 92121 (US). (72) Inventors: HOWELL, Mark, D. ; 8846 Padoga Way, San Diego, CA 92126 (US). BROSTOFF, Steven, W. ; 2608 La Golondrina Street, Carlsbad, CA 92008 (US). CARLO, Dennis, J. ; 4466 Los Pinos, Rancho Santa Fe, CA 92067 (US).		326,314	21 March 1989 (21.03.89)	US	382,085	18 July 1989 (18.07.89)	US	382,086	18 July 1989 (18.07.89)	US	(74) Agents: CAMPBELL, Cathryn et al.; Pretty, Schroeder, Brueggemann & Clark, 444 South Flower Street, Suite 2000, Los Angeles, CA 90071 (US). (81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), + CH (European patent), CM (OAPI patent), DE (European patent), DK, DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
326,314	21 March 1989 (21.03.89)	US									
382,085	18 July 1989 (18.07.89)	US									
382,086	18 July 1989 (18.07.89)	US									
(54) Title: VACCINATION AND METHODS AGAINST DISEASES RESULTING FROM PATHOGENIC RESPONSES BY SPECIFIC T CELL POPULATIONS											
(57) Abstract <p>The present invention provides vaccines and a means of vaccinating a mammal so as to prevent or control specific T cell mediated pathologies or to treat the unregulated replication of T cells. The vaccine is composed of a T cell receptor (TCR) or a fragment thereof corresponding to a TCR present on the surface of T cells mediating the pathology. The vaccine fragment can be a peptide corresponding to sequences of TCRs characteristic of the T cells mediating said pathology. Means of determining appropriate amino acid sequences for such vaccines are also provided. The vaccine is administered to the mammal in a manner that induces an immune response directed against the TCR of T cells mediating the pathology. This immune response down regulates or deletes the pathogenic T cells, thus ablating the disease pathogenesis. The invention additionally provides a specific β-chain variable region of the T cell receptor, designated Vβ17, which is central to the pathogenesis of rheumatoid arthritis (RA). Also provided are means to detect, prevent and treat RA.</p>											

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

VACCINATION AND METHODS AGAINST DISEASES RESULTING
FROM PATHOGENIC RESPONSES
BY SPECIFIC T CELL POPULATIONS

BACKGROUND OF THE INVENTION

5 This invention relates to the immune system and, more specifically, to methods of modifying pathological immune responses.

Higher organisms are characterized by an immune system which protects them against invasion by potentially deleterious substances or microorganisms. When a substance, termed an antigen, enters the body, and is recognized as foreign, the immune system mounts both an antibody-mediated response and a cell-mediated response. Cells of the immune system termed B lymphocytes, or B cells, produce antibodies which specifically recognize and bind to the foreign substance. Other lymphocytes termed T lymphocytes, or T cells, both effect and regulate the cell-mediated response resulting eventually in the elimination of the antigen.

20 A variety of T cells are involved in the cell-mediated response. Some induce particular B cell clones to proliferate and produce antibodies specific for the antigen. Others recognize and destroy cells presenting foreign antigens on their surfaces. Certain T cells regulate the response by either stimulating or suppressing other cells.

While the normal immune system is closely regulated, aberrations in immune response are not uncommon. In some instances, the immune system functions inappropriately and reacts to a component of the host as if it were, in fact, foreign. Such a response results in an autoimmune disease, in which the host's immune system attacks the host's own tissue. T cells, as the primary regulators of the immune

system, directly or indirectly effect such autoimmune pathologies.

Numerous diseases are believed to result from autoimmune mechanisms. Prominent among these are
5 rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, Type I diabetes, myasthenia gravis and pemphigus vulgaris. Autoimmune diseases affect millions of individuals world-wide and the cost of these diseases, in terms of actual treatment and expenditures and lost
10 productivity, is measured in billions of dollars annually. At present, there are no known effective treatments for such autoimmune pathologies. Usually, only the symptoms can be treated, while the disease continues to progress, often resulting in severe debilitation or death.

15 In other instances, lymphocytes replicate inappropriately and without control. Such replication results in a cancerous condition known as a lymphoma. Where the unregulated lymphocytes are of the T cell type, the tumors are termed T cell lymphomas. As with other
20 malignancies, T cell lymphomas are difficult to treat effectively.

Thus there exists a long-felt need for an effective means of curing or ameliorating T cell mediated pathologies. Such a treatment should ideally control the
25 inappropriate T cell response, rather than merely reducing the symptoms. The present invention satisfies this need and provides related advantages as well.

Summary of the Invention

The present invention provides vaccines and a means of vaccinating a mammal so as to prevent or control specific T cell mediated pathologies or to treat the unregulated
5 clonal replication of T cells. The vaccine is composed of a T cell receptor (TCR) or a fragment thereof corresponding to a TCR present on the surface of T cells mediating the pathology. The vaccine fragment can be a peptide corresponding to sequences of TCRs characteristic of the T
10 cells mediating said pathology.

Means of determining appropriate amino acid sequences for such vaccines are also provided. The vaccine is administered to the mammal in a manner that induces an immune response directed against the TCR of T cells
15 mediating the pathology. This immune response down regulates or deletes the pathogenic T cells, thus ablating the disease pathogenesis.

The invention additionally provides a specific β -chain variable region of the T cell receptor, designated
20 VB17, which is central to the pathogenesis of rheumatoid arthritis (RA). Also provided are means to detect, prevent and treat RA.

The invention also provides a specific region of a T cell receptor useful for the treatment of multiple
25 sclerosis (MS). Also provided are means to detect, prevent and treat MS.

Detailed Description of the Invention

The invention relates to vaccines and their use for preventing or ameliorating T cell-mediated pathologies,
30 such as autoimmune diseases and T cell lymphomas. Vaccination provides a specific and sustained treatment

which avoids problems associated with other potential avenues of therapy.

As used herein, the term "T cell-mediated pathology" refers to any condition in which an inappropriate T cell response is a component of the pathology. The term is intended to include both diseases directly mediated by T cells and those, such as myasthenia gravis, which are characterized primarily by damage resulting from antibody binding, and also diseases in which an inappropriate T cell response contributes to the production of those antibodies. The term is intended to encompass both T cell mediated autoimmune diseases and unregulated clonal T cell replication.

As used herein, "substantially the sequence" when referring to an amino acid sequence means the described sequence or other sequences having any additions, deletions or substitutions which do not substantially effect the ability of the sequence to elicit an immune response against the desired T cell receptor sequence. Thus, a portion of the described immunizing sequence can be used so long as it is sufficiently characteristic of the desired T cell receptor as to cause an effective immune response against desired T cell receptors but not against undesired T cell receptors. Such variations in the sequence can easily be made, e.g. by synthesizing an alternative sequence, and tested, e.g. by immunizing a mammal, to determine its effectiveness.

As used herein, the term "fragment" is intended to cover such fragments in conjunction with or combined with additional sequences or moieties, as for example where the peptide is coupled to other amino acid sequences or to a carrier. The terms "fragment" and "peptide" can, therefore, be used interchangeably since a peptide will be the most common fragment of the T cell receptor. Each

fragment of the invention can have an altered sequence, as described above for the term "substantially the sequence."

Reference herein to a "fragment or portion of the T cell receptor" does not mean that the composition must be derived from intact T cell receptors. Such "fragments or portions" can be produced by various means well-known to those skilled in the art, such as for example manual or automatic peptide synthesis or methods of cloning.

As used herein when referring to the relationship between peptide fragments of the invention and sequences of TCRs, "corresponding to" means that the peptide fragment has an amino acid sequence which is sufficiently homologous to the TCR sequence to stimulate an effective regulatory response in the individual. The sequence need not be identical to the TCR sequence, however, as shown in Examples II and III.

By "immunogenically effective" is meant an amount of the T cell receptor or fragment thereof which, is effective to elicit an immune response to prevent or treat a T cell mediated pathology or an unregulated T cell clonal replication in the individual. Obviously, such amounts will vary between species and individuals depending on many factors. For example, higher doses will generally be required for an effective immune response in a human compared with a mouse.

As used herein, "VB17" refers to a specific B-chain variable region of a T cell receptor (TCR). VB17 has the amino acid sequence MSNQVLCCVVLCLGANTVDGGITQSPKYLFRKEGQN VTLSC EQNLNHDAMYWYRQDPGQGLRLIYYSQIVNDFQKGDIAEGYSVSREKKESFP LTVTSAQKNPTAFYLCASS. The hypervariable and junctional regions are most useful for vaccines. One hypervariable region of VB17 especially useful is the CDR2 region which has the amino acid sequence SQIVNDFQK. Modifications in

this sequence which do not affect the ability of the receptor to act as an immunogen to stimulate the desired immune response are also included in the definition. The variable region can be joined with any D and J segment of the TCR. Further, immunogenically representative fragments of VB17 are also included in the definition of "VB17."

As used herein, "binding partner" means a compound which is reactive with a TCR. Generally, this compound will be a Major Histocompatibility Antigen (MHC) but can be any compound so long as when the TCR is bound in the normal course, T cell activation or proliferation occurs.

As used herein, "ligand" means any molecule that reacts to form a complex with another molecule.

As used herein, "selectively binds" means that a molecule binds to one type of molecule but not substantially to other types of molecules. In relation to VB17 "selective binding" indicates binding to VB17 containing TCRs but not substantially to other TCRs which lack VB17.

The immune system is the primary biological defense of the host (self) against potentially pernicious agents (non-self). These pernicious agents may be pathogens, such as bacteria or viruses, as well as modified self cells, including virus-infected cells, tumor cells or other abnormal cells of the host. Collectively, these targets of the immune system are referred to as antigens. The recognition of antigen by the immune system rapidly mobilizes immune mechanisms to destroy that antigen, thus preserving the sanctity of the host environment.

The principal manifestations of an antigen-specific immune response are humoral immunity (antibody mediated) and cellular immunity (cell mediated). Each of these

immunological mechanisms are initiated through the activation of helper (CD4+) T Cells. These CD4+ T cells in turn stimulate B cells, primed for antibody synthesis by antigen binding, to proliferate and secrete antibody. This
5 secreted antibody binds to the antigen and facilitates its destruction by other immune mechanisms. Similarly, CD4+ T cells provide stimulatory signals to cytotoxic (CD8+) T cells which recognize and destroy cellular targets (for example, virus infected cells of the host). Thus, the
10 activation of CD4+ T cells is the proximal event in the stimulation of an immune response. Therefore, elaboration of the mechanisms underlying antigen specific activation of CD4+ T cells is crucial in any attempt to selectively modify immunological function.

15 T cells owe their antigen specificity to the T cell receptor (TCR) which is expressed on the cell surface. The TCR is a heterodimeric glycoprotein, composed of two polypeptide chains, each with a molecular weight of approximately 45 kD. Two forms of the TCR have been
20 identified. One is composed of an alpha chain and a beta chain, while the second consists of a gamma chain and a delta chain. Each of these four TCR polypeptide chains is encoded by a distinct genetic locus containing multiple discontinuous gene segments. These include variable (V)
25 region gene segments, junction (J) region gene segments and constant (C) region gene segments. Beta and delta chains contain an additional element termed the diversity (D) gene segment. (Since D segments and elements are found in only some of the TCR genetic loci, and polypeptides, further
30 references herein to D segments and elements will be in parentheses to indicate the inclusion of these regions only in the appropriate TCR chains. Thus, V(D)J refers either to VDJ sequences of chains which have a D region or refers to VJ sequences of chains lacking D regions.)

35 During lymphocyte maturation, single V, (D) and J gene

segments are rearranged to form a functional gene that determines the amino acid sequence of the TCR expressed by that cell. Since the pool of V, (D) and J genes which may be rearranged is multi-membered and since individual
5 members of these pools may be rearranged in virtually any combination, the complete TCR repertoire is highly diverse and capable of specifically recognizing and binding the vast array of binding partners to which an organism may be exposed. However, a particular T cell will have only one
10 TCR molecule and that TCR molecule, to a large degree if not singly, determines the specificity of that T cell for its binding partner.

Animal models have contributed significantly to our
15 understanding of the immunological mechanisms of autoimmune disease. One such animal model, experimental allergic encephalomyelitis (EAE), is an autoimmune disease of the central nervous system that can be induced in mice and rats by immunization with myelin basic protein (MBP). The
20 disease is characterized clinically by paralysis and mild wasting and histologically by a perivascular mononuclear cell infiltration of the central nervous system parenchyma. The disease pathogenesis is mediated by T cells with specificity for MBP. Multiple clones of MBP-specific T
25 cells have been isolated from animals suffering from EAE and have been propagated in continuous culture. After in vitro stimulation with MBP, these T cell clones rapidly induce EAE when adoptively transferred to healthy hosts. Importantly, these EAE-inducing T cells are specific, not
30 only for the same antigen (MBP), but also usually for a single epitope on that antigen. These observations indicate that discrete populations of autoaggressive T cells are responsible for the pathogenesis of EAE.

Analysis of the TCRs of EAE-inducing T cells has
35 revealed restricted heterogeneity in the structure of these disease-associated receptors. In one analysis of 33 MBP-

reactive T cells, only two alpha chain V region gene segments and a single alpha chain J region gene segment were utilized. Similar restriction of beta chain TCR gene usage was also observed in this T cell population. Only
5 two beta chain V region segments and two J region gene segments were found. More importantly, approximately eighty percent of the T cell clones had identical amino acid sequences across the region of beta chain V-D-J joining. These findings confirm the notion of common TCR
10 structure among T cells with similar antigen specificities and indicate that the TCR is an effective target for immunotherapeutic strategies aimed at eliminating the pathogenesis of EAE.

Various attempts have been made to exploit the
15 antigen specificity of autoaggressive T cells in devising treatment strategies for EAE. For example, passive administration of monoclonal antibodies specific for TCRs present on EAE-inducing T cells has been employed. In the mouse model of EAE, infusion of a monoclonal antibody
20 specific for V_β8, the major beta chain V region gene used by MBP-specific T cells, reduced the susceptibility of mice to subsequent EAE induction (Acha-Orbea et al., Cell 54:263-273 (1988) and Urban et al., Cell 54:577-592 (1988)). Similar protection has been demonstrated in rat EAE with
25 monoclonal antibody reactive with an unidentified idiotypic determinant of the TCR on MBP specific T cells (Burns et al., J. Exp. Med. 169:27-39 (1989)). While passive antibody therapy appears to have some ameliorative effect on EAE susceptibility, it is fraught with potential
30 problems. The protection afforded is transient, thus requiring repeated administration of the antibody. Multiple infusions of antibody increases the chances that the host will mount an immune response to the administered antibody, particularly if it is raised in a xenogeneic
35 animal. Further an antibody response to a pathogenic T cell clone represents only one element in the complete

immune response and neglects the potential contributions of cellular immunity in resolving the autoreactivity.

The role of cellular immunity in reducing the activity of autoaggressive T cells in EAE has been examined and potential therapies suggested. In a manner similar to the passive antibody approach, regulatory T cells have been derived ex vivo and readministered for immunotherapy. For example, Sun et al., Nature, 332:843-845 (1988), have recently isolated a CD8+ T cell clone from convalescing rats in whom EAE had been induced by adoptive transfer of an MBP-specific CD4+ T cell line. This CD8+ T cell clone displayed cytolytic activity in vitro for the CD4+ T cell used to induce disease. Moreover, adoptive transfer of this CTL clone reduced the susceptibility of recipient rats to subsequent challenge with MBP. Lider et al., Science, 239:181-183 (1988) have also isolated a CD8+ T cell clones with suppressive activity for EAE-inducing T cells. This CD8+ clone was isolated from rats vaccinated with attenuated disease-inducing T cell clones and, though it showed no cytolytic activity in vitro, it could suppress MBP-driven proliferation of EAE-inducing T cells. Although these studies indicate that the CD8+ T cells could downregulate EAE, it is hard to reconcile a major role for these selected CD8+ CTLs in the long-term resistance of the recovered rats since Sedgwick, et al., (Eur. J. Immunol., 18:495-502 (1988)) have clearly shown that depletion of CD8+ cells with monoclonal antibodies does not affect the disease process or recovery.

In the experiments of Sun et al., and Lider et al., described above, the administration of extant derived regulatory T cells overcomes the major obstacle of passive antibody therapy; it permits a regulatory response in vivo of prolonged duration. However, it requires in vitro cultivation with attenuated disease-inducing T cells to develop clones of such regulatory T cells, a costly and

labor intensive process. Further, in an outbred population such as humans, MHC non-identity among individuals makes this a highly individualized therapeutic strategy. Regulatory clones need to be derived for each individual patient and then re-administered only to that patient to avoid potential graft versus host reactions.

Direct vaccination with attenuated disease-inducing T cell clones also has been employed as a therapy for EAE. MBP-specific T cells, capable of transferring disease, have been attenuated by gamma irradiation or chemical fixation and used to vaccinate naive rats. In some cases, vaccinated animals exhibited resistance to subsequent attempts at EAE induction (Lider et al., supra; see Cohen and Weiner, Immunol. Today 9:332-335 (1988) for review). The effectiveness of such vaccination, however, is inconsistent and the degree of protection is highly variable. T cells contain a multitude of different antigens which induce an immune response when the whole T cell is administered as a vaccine. This phenomenon has been demonstrated by Offner et al., (J. Neuroimmunol., 21:13-22 (1989)), who showed that immunization with whole T cells increased the delayed type hypersensitivity (DTH) response, as measured by ear swelling, to those T cells in an incremental manner as the number of vaccinations increased. However, positive DTH responses were found in both protected and non-protected animals. Rats responded similarly to both the vaccinating encephalitogenic T cells and control T cells. Conversely, vaccination with PPD-specific T cells from a PPD-specific T cell line induced DTH to the vaccinating cells as well as to an encephalitogenic clone even though no protection was observed. The similar response of vaccinated rats to both disease-inducing and control cells, as quantified by delayed-type hypersensitivity (a measure of cell-mediated immunity), indicates that numerous antigens on these T cells are inducing immune responses. Thus, vaccination

with attenuated disease-inducing T cells suffers from a lack of specificity for the protective antigen on the surface of that T cell, as well as, variable induction of immunity to that antigen. As a candidate for the treatment
5 of human diseases, vaccination with attenuated T cells is plagued by the same labor intensiveness and need for individualized therapies as noted above for infusion of CD8+ cells.

The present invention provides an effective method of
10 immunotherapy for T cell mediated pathologies, including autoimmune diseases, which avoids many of the problems associated with the previously suggested methods of treatment. By vaccinating, rather than passively administering heterologous antibodies, the host's own
15 immune system is mobilized to suppress the autoaggressive T cells. Thus, the suppression is persistent and may involve any and all immunological mechanisms in effecting that suppression. This multi-faceted response is more effective than the uni-dimensional suppression achieved by
20 passive administration of monoclonal antibodies or extant-derived regulatory T cell clones.

As they relate to autoimmune disease, the vaccines of the present invention comprise TCRs of T cells that mediate autoimmune diseases. The vaccines can be whole TCRs
25 substantially purified from T cell clones, individual T cell receptor chains (for example, alpha, beta, etc.) or portions of such chains, either alone or in combination. The vaccine can be homogenous, for example, a single peptide, or can be composed of more than one type of
30 peptide, each of which corresponds to a different portion of the TCR. Further, these peptides can be from distinct TCRs wherein both TCRs contribute to the T cell mediated pathology.

In a specific embodiment, the immunizing peptide can

have the amino acid sequence SGDQGGNE when the subject has multiple sclerosis. Any immunogenic portion of this peptide can be effective. Thus, amino acid substitutions can be made which do not destroy the immunogenicity of the peptide. Additionally, this peptide can be linked to a carrier to further increase its immunogenicity.

In a further specific embodiment, T cell receptors or fragments of the TCR which contain VB17 can be used to immunize an individual having rheumatoid arthritis to treat or prevent the disease. The immune response generated in the individual can neutralize or kill T cells having VB17 and, thus, prevent or treat the deleterious effects of VB17-bearing T cells. Moreover, to the extent that VB17 is common to T cell receptors on pathogenic T cells mediating autoimmune diseases in general, such vaccines can also be effective in ameliorating such other autoimmune diseases.

By "substantially pure" it is meant that the TCR is substantially free of other biochemical moieties with which it is normally associated in nature. Alternatively, the vaccines comprise peptides of varying lengths corresponding to the TCR or portions thereof. The peptides can be produced synthetically or recombinantly, by means well known to those skilled in the art. Preferably, the peptide vaccines correspond to regions of the TCR which distinguish that TCR from other nonpathogenic TCRs. Such specific regions can be located within the various region(s) of the respective TCR polypeptide chains, especially a short sequence spanning the V(D)J junction, thus restricting the immune response solely to those T cells bearing this single determinant.

The vaccines are administered to a host exhibiting or at risk of exhibiting an autoimmune response. Definite clinical diagnosis of a particular autoimmune disease

warrants the administration of the relevant disease-specific TCR vaccines. Prophylactic applications are warranted in diseases where the autoimmune mechanisms precede the onset of overt clinical disease (for example, 5 Type I Diabetes). Thus, individuals with familial history of disease and predicted to be at risk by reliable prognostic indicators could be treated prophylactically to interdict autoimmune mechanisms prior to their onset.

TCR vaccines can be administered in many possible 10 formulations, in pharmacologically acceptable mediums. In the case of a short peptide, the peptide can be conjugated to a carrier, such as KLH, in order to increase its immunogenicity. The vaccine can be administered in conjunction with an adjuvant, various of which are known to 15 those skilled in the art. After initial immunization with the vaccine, a booster can be provided. The vaccines are administered by conventional methods, in dosages which are sufficient to elicit an immunological response, which can be easily determined by those skilled in the art.

20 Appropriate peptides to be used for immunization can be determined as follows. Disease-inducing T cell clones reactive with the target antigens are isolated from affected individuals. Such T cells are obtained preferably from the site of active autoaggressive activity such as a 25 lesion in the case of pemphigus vulgaris, central nervous system (CNS) in the case of multiple sclerosis or synovial fluid or tissue in the case of rheumatoid arthritis, or alternatively from blood of affected individuals. The TCR genes from these autoaggressive T cells are then sequenced. 30 Polypeptides corresponding to TCRs or portions thereof that are selectively represented among disease inducing T cells (relative to non-pathogenic T cells) can then be selected as vaccines and made and used as described above.

Alternatively, the vaccines can comprise anti-

idiotypic antibodies which are internal images of the peptides described above. Methods of making, selecting and administering such anti-idiotypic vaccines are well known in the art. See, for example, Eichmann, et al., CRC Critical
5 Reviews in Immunology 7:193-227 (1987), which is incorporated herein by reference.

T Cell Pathologies of Malignant Etiology

To illustrate the utility of TCR vaccination, autoimmune disease has been discussed. However, T cell
10 lymphoma is another T cell pathology which would be amenable to this type of treatment. Application of this technology in the treatment of T lymphoma would be conducted in virtually identical fashion. In one respect, however, this technology is more readily applied to T cell
15 proliferative disease since the isolation of the pathogenic T cells is more easily accomplished. Once the clones are isolated, the technology is applied in the manner described herein. Specifically, the TCR genes of the T lymphomas are sequenced, appropriate regions of those TCRs are identified
20 and used as vaccines. The vaccines can comprise single or multiple peptides, and can be administered in pharmacologically acceptable formulations with or without adjuvants by conventional means.

Multiple Sclerosis

25 T cells causative of multiple sclerosis (MS) have not previously been identified, though MBP-reactive T cells have been proposed to play a role due to the clinical and histologic similarities between MS and EAE. In rat and mouse models of EAE, MBP-reactive, encephalogenic T cells
30 show striking conservation of β -chain VDJ amino acid sequence, despite known differences in MHC restriction and MBP-peptide antigen specificity. This invention is premised on the observation that a human myelin basic

protein (MBP)-reactive T cell line, derived from an MS patient, has a TCR β -chain with a VDJ amino acid sequence homologous with that of β -chains from MBP-reactive T cells mediating pathogenesis in experimental allergic encephalomyelitis (EAE), an animal model of MS. This line is specific for another epitope of MBP. This finding demonstrates the involvement of MBP-reactive T cells in the pathogenesis of MS and demonstrates that TCR peptides similar to those described herein for the prevention of EAE can be appropriate in treating MS.

Rheumatoid Arthritis

Rheumatoid arthritis (RA) is a T cell mediated autoimmune disease. The invention describes oligoclonal infiltrates of activated VB17 T cells in the synovium of rheumatoid arthritis patients. The presence of these T cells in the diseased tissue of all patients examined, their oligoclonality, and the cytotoxic activity of one such T cell for synovial adherent cells, demonstrates a central role for VB17 bearing T cells in the pathogenesis of RA.

Activated T cell populations in the synovial tissue of RA patients have been examined by analyzing T cell receptor (TCR) mRNAs isolated from IL-2 receptor positive (IL-2R+) synovial T cells. TCR mRNAs were amplified using a polymerase chain reaction (PCR) protocol designed to amplify human TCR β -chain genes containing virtually any VB gene element. In this analysis, oligoclonal VB17 rearrangements were found to be enriched in the IL2-R+ population, indicating that VB17 T cells are likely involved in the pathogenesis of RA. A CD4+, VB17 bearing T cell clone has been isolated from one of the synovial tissue specimens and its in vitro cytotoxicity for synovial adherent cells supports the direct involvement of VB17 T cells in RA.

As noted, the invention provides the extremely important discovery that a specific variable region of the β -chain of the TCR, designated VB17, is closely associated with rheumatoid arthritis in human subjects. This
5 discovery allows for the detection, prevention and treatment of rheumatoid arthritis using the methodology set out in this invention. Similar therapeutic approaches set out above for EAE can be applied to rheumatoid arthritis by those skilled in the art.

10 Specifically, the invention provides a method of diagnosing or predicting susceptibility to rheumatoid arthritis in an individual comprising detecting T cells having the β -chain variable region designated VB17 in a sample from the individual, the presence abnormal levels of
15 VB17-containing T cells indicating rheumatoid arthritis or susceptibility to rheumatoid arthritis. The VB17 containing T cell can be qualitatively or quantitatively compared to that of a normal individual. Such diagnosis can be performed by detecting a portion of the VB17 which
20 does not occur on non-rheumatoid arthritis associated β -chain variable region T-cell receptors. The VB17 can be detected, for example, by contacting the VB17 with a detectable ligand capable of specifically binding to VB17. Many such detectable ligands are known in the art, e.g. an
25 enzyme linked antibody. Alternatively, nucleotide probes complementary to VB17 encoding nucleic acid sequences can be utilized to detect VB17 containing T cells, as taught in Example IX.

The invention also provides a method of preventing or
30 treating rheumatoid arthritis comprising preventing the attachment of a VB17 containing T-cell receptor to its binding partner. In one embodiment attachment is prevented by binding a ligand to VB17. In an alternative embodiment attachment is prevented by binding a ligand to the VB17

binding partner. Attachment can be prevented by known methods, e.g. binding an antibody to VB17 or the binding portion to physically block attachment.

The invention also provides a method of preventing or
5 treating rheumatoid arthritis in an individual comprising cytotoxicly or cytostatically treating VB17 containing T-cells in the individual. In one embodiment, the VB17 containing T-cells are treated with a cytotoxic or cytostatic agent which selectively binds VB17. The agent
10 can be an antibody attached to a radioactive or chemotherapeutic moiety. Such attachment and effective agents are well known in the art. See, for example, Harlow, E. and Lane, Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, 1988, which is incorporated
15 herein by reference.

The invention also provides the extremely important discovery that a specific TCR sequence, SGDQGGNE, is closely associated with multiple sclerosis in human subjects. This discovery allows for the detection,
20 prevention and treatment of multiple sclerosis using the methodology set out in this invention. Similar therapeutic approaches set out herein for EAE can be applied to multiple sclerosis by those skilled in the art.

Specifically, the invention provides a method of
25 diagnosing or predicting susceptibility to multiple sclerosis in an individual comprising detecting T cells having substantially the SGDQGGNE sequence in a sample from the individual, the presence of the sequence indicating multiple sclerosis or susceptibility to multiple sclerosis.
30 The sequence can be detected, for example, by contacting it with a detectable ligand. Many such ligands are known in the art, e.g. an enzyme linked antibody. Alternatively, nucleotide probes complementary to the nucleic acid encoding the sequence can be utilized to detect T cells as,

taught in Example IX.

The invention also provides a method of preventing or treating multiple sclerosis comprising preventing the attachment of a T-cell receptor containing substantially the SGDQGGNE sequence to its binding partner. In one embodiment attachment is prevented by binding a ligand to to the sequence. In an alternative embodiment attachment is prevented by binding a ligand to the binding partner. Attachment can be prevented by known methods, e.g. binding an antibody to the sequence to physically block attachment.

The invention also provides a method of preventing or treating multiple sclerosis in an individual comprising cytotoxicly or cytostatically treating T cells containing substantially the SGDQGGNE sequence in the individual. In one embodiment, T-cells are treated with a cytotoxic or cytostatic agent which selectively binds the sequence. The agent can be an antibody attached to a radioactive or chemotherapeutic moiety.

The following examples are intended to illustrate but not limit the invention.

EXAMPLE I RAT MODEL OF EAE

Female Lewis rats, (Charles River Laboratories, Raleigh-Durham, NC) were immunized in each hind foot pad with 50 μ g of guinea pig myelin basic protein emulsified in complete Freund's adjuvant. The first signs of disease were typically observed 9-11 days post-immunization. Disease severity is scored on a three point scale as follows: 1=limp tail; 2=hind leg weakness; 3=hind leg paralysis. Following a disease course of approximately four to six days, most rats spontaneously recovered and were refractory to subsequent EAE induction.

EXAMPLE II
SELECTION AND PREPARATION OF VACCINES

Vaccinations were conducted with a T cell receptor peptide whose sequence was deduced from the DNA sequence of a T cell receptor beta gene predominating among EAE-inducing T cells of BIO.PL/L mice. The DNA sequence was that reported by Urban, et al., supra, which is incorporated herein by reference. A nine amino acid peptide, having the sequence of the VDJ junction of the TCR beta chain of the mouse, was synthesized by methods known to those skilled in the art. The sequence of this peptide is: SGDAGGGYE. (Amino acids are represented by the conventional single letter codes.) The equivalent sequence in the rat has been reported to be: SSD-SSNTE (Burns et al., J. Exp. Med. 169:27-39 (1989)). The peptide was desalted by Sephadex G-25 (Pharmacia Fine Chemicals, Piscataway, NJ) column chromatography in 0.1 M acetic acid and the solvent was subsequently removed by two cycles of lyophilization. A portion of the peptide was conjugated to keyhole limpet hemocyanin (KLH) with glutaraldehyde at a ratio of 7.5 mgs of peptide per mg of KLH. The resulting conjugate was dialyzed against phosphate buffered saline (PBS).

EXAMPLE III
VACCINATION AGAINST EAE

Vaccines used in these studies consisted of free VDJ peptide and also of VDJ peptide conjugated to KLH. These were dissolved in PBS and were emulsified with equal volumes of either (1) incomplete Freund's adjuvant (IFA) or (2) complete Freund's adjuvant (CFA) made by suspending 10 mg/ml heat killed desiccated Mycobacterium tuberculosis H37ra (Difco Laboratories, Detroit, MI) in IFA. Emulsions

15

TABLE I

40

EXAMPLE IVVaccination against EAE with Lewis Rat VDJ peptides

The VDJ peptide used in the previous examples was synthesized according to the sequence of TCR β chain molecules found on EAE-inducing T cells in B10.PL mice. In addition, peptides were synthesized and tested which correspond to sequences found on encephalitogenic T cells in Lewis rats. These VDJ sequences are homologous with that of B10.PL mice, but not identical. The rat peptides were synthesized according to the DNA sequences reported by Burns, et al. and Chluba, et al., Eur. J. Immunol. 19:279-284 (1989). The sequences of these three peptides designated IR1, 2 and 3, are shown below, aligned with the B10.PL mouse sequence used in Examples I through III (VDJ).

15	VDJ	S G D A G G Y E
	IR1	C A S S D - S S N T E V F F G K
	IR2	C A S S D - S G N T E V F F G K
	IR3	C A S S D - S G N - V L Y F G E G S R
	IR9b	A S S D - S S N T E

20 The preparation, administration and evaluation of these vaccines were conducted as described in Examples I through III with the following exceptions: 50 μ g of the individual VDJ peptides were incorporated into vaccine formulations containing CFA; neither vaccinations in IFA nor vaccinations with peptides conjugated to KLH were conducted. Control animals were untreated prior to MBP challenge as in Example III or were vaccinated with emulsions of PBS and CFA to assess the protective effect of adjuvant alone. The results are shown in Table II below.

TABLE II

	Animal No.	Vaccination (Adjuvant)	Days After Challenge								
			10	11	12	13	14	15	16	17	18
5	1	None	-	1	2	3	3	2	-	-	-
	2	"	1	3	3	3	2	-	-	-	-
	3	"	-	2	3	3	3	1	-	-	-
10	4	PBS-CFA	1	2	3	3	3	-	-	-	-
	5	"	1	2	3	3	3	-	-	-	-
	6	"	-	2	3	3	3	-	-	-	-
	7	IR1 (50 µg)	-	-	-	2	1	-	-	-	-
	8	"	-	-	-	-	1	3	-	-	-
	9	"	-	-	-	1	1	1	1	-	-
15	10	IR2 (50µg)	-	-	1	3	3	3	-	-	-
	11	"	-	-	-	-	2	2	3	3	-
	12	"	-	-	-	-	1	-	-	-	-
	13	IR3 (50µg)	1	3	3	3	2	-	-	-	-
	14	"	-	-	2	3	3	-	-	-	-
	15	"	-	-	-	-	-	-	-	-	-
20	16	IR9b (50 µg)	-	-	-	-	-	-	-	-	-
	17	"	-	-	-	-	-	-	-	-	-
	18	"	-	-	-	-	-	-	-	-	-
	19	"	-	-	-	-	-	-	-	-	-
25	Scoring: - no signs										
		1)	limp tail								
		2)	hind leg weakness								
		3)	hind leg paralysis								

As shown in Table II, disease in unvaccinated control animals was observed as early as day 10. Disease was characterized by severe paralysis and wasting, persisted for 4 to 6 days and spontaneously remitted. PBS-CFA vaccinated rats displayed disease courses virtually indistinguishable from those of unvaccinated controls. In contrast, delays in onset were observed in some of the IR1, 2 or 3 vaccinated animals and others showed both delayed onset as well as decreased severity and/or duration of disease. Overall, however, vaccinations with the rat VDJ peptides (IR1-3) were slightly less effective than those

with the mouse VDJ peptide (Example III). Vaccination with IR9b, however, afforded complete protection in all four animals in which it was tested. Importantly, no histologic lesions characteristic of disease were found in any of the 5 four animals vaccinated with IR9b indicating that sub-clinical signs of disease were also abrogated.

EXAMPLE V

Vaccination with V region specific peptides

A peptide specific for the V β 8 gene family was tested 10 as a vaccine against EAE. V β 8 is the most common β chain gene family used by encephalitogenic T cells in both rats and mice. A peptide was synthesized based on a unique DNA sequence found in the V β 8 gene, and which is not found among other rat V β genes whose sequences were reported by 15 Morris, et al., Immunogenetics 27:174-179 (1988). The sequence of this V β 8 peptide, designated IR7, is:

IR7 D M G H G L R L I H Y S Y D V N S T E K

The efficacy of this V β 8 peptide was tested in the Lewis rat model of EAE (Example I) as described in Examples 20 II and III. 50 μ g of peptide were tested in CFA. Vaccinations in IFA or with peptide-KLH conjugates were not conducted. The results of these studies are shown in Table III.

25

TABLE III

	Animal No.	Vaccination (Adjuvant)	Days After Challenge								
			10	11	12	13	14	15	16	17	18
5	1	IR7 (50 µg)	-	-	1	2	3	3	3	-	-
	2	"	-	-	-	-	1	1	-	-	-
	3	"	-	-	-	-	-	-	-	-	-
10	Scoring:	-	no signs								
		1)	limp tail								
		2)	hind leg weakness								
		3)	hind leg paralysis								

The results of vaccinations conducted with the rat V88 peptide are similar to those observed with the mouse and rat IR1, 2 and 3 peptides. Delayed onset as well as decreased severity and duration of disease was observed in one animal. One animal was completely protected.

EXAMPLE VIVaccination with J region peptides

A peptide was synthesized which corresponds to the J α gene segment, TA39, found among both rat and mouse encephalitogenic T cell receptors. The sequence of this peptide, designated IR5, is:

IR5 R F G A G T R L T V K

The efficacy of the J α TA39 peptide was tested in the Lewis rat model of EAE (Example I) as described in Examples II and III. 50 µg of peptide were tested in CFA. Vaccinations in IFA or with peptide-KLH conjugates were not conducted. The results of these studies are shown in Table IV.

TABLE IV

Animal No.	Vaccination (Adjuvant)	Days After Challenge											
		10	11	12	13	14	15	16	17	18	19	20	
5	1	IR5 (50 μ g)	-	-	-	-	-	2	1	1	1	1	-
	2	"	-	-	-	-	-	-	-	-	-	-	-
	3	"	-	-	-	-	-	-	-	-	-	-	-
<hr/>													
10	Scoring:	-	no signs										
		1)	limp tail										
		2)	hind leg weakness										
		3)	hind leg paralysis										

The results of vaccinations conducted with the rat J α TA39 peptide are more effective than those observed with the mouse VDJ peptide or the VB8 peptide. Two of three animals were totally protected and, in the third, disease onset was markedly delayed. Severity was also reduced in this animal though disease persisted for a normal course of 5 days. Importantly, the two animals which were completely protected showed no histologic evidence of T cell infiltration of the CNS. This result indicates that vaccinating with the J α TA39 very efficiently induces a regulatory response directed at encephalitogenic T cells. Even sub-clinical signs of disease were abrogated.

25

EXAMPLE VIIVaccination with mixtures of TCR peptides

Vaccinations were conducted with a mixture of TCR peptides. This mixture contained 50 μ g of each of the peptides IR1, 2, 3 and 5 (the three rat VDJ peptides and the rat J α TA39 peptide).

The efficacy of this peptide mixture was tested in the

Lewis rat model (Example I) as described in Examples II and III. Peptides were tested in CFA. Vaccinations in IFA or with peptide-KLH conjugates were not conducted. The results of these studies are shown in Table V.

5

TABLE V

	Animal No.	Vaccination (Adjuvant)	Days After Challenge									
			10	11	12	13	14	15	16	17	18	
10	4	IR1, 2, 3, 5	-	-	-	-	-	-	-	-	-	
	5	(50 µg each)	-	-	-	-	-	-	-	-	-	
	6	"	-	-	-	-	-	-	-	-	-	
15	Scoring:	-	no signs									
		1)	limp tail									
		2)	hind leg weakness									
		3)	hind leg paralysis									

The results of vaccinations conducted with the rat JαTA39 and three VDJ peptides are almost as effective as those described for IR9b in Table II. All three animals were totally protected. In addition to the absence of any clinical signs of EAE, two of these three animals were completely free of histological evidence of T cell infiltration into the CNS while the third showed only two small foci of lymphocytic infiltration at the base of the spinal cord.

EXAMPLE VIII

Multiple Sclerosis Vaccine

Human MBP-reactive T cells

MBP-reactive T cell lines were established from peripheral blood mononuclear cells (PBMC) of nine chronic progressive MS patients and two healthy controls. Cells were maintained in culture by regular stimulation with

purified human MBP and irradiated-autologous PBMC for three days followed by four days in IL-2 containing medium.

PCR Amplification of TCR β -chain genes from MBP-reactive T cell lines

- 5 T cells were harvested from log phase cultures and RNA was prepared, amplified with the VB16mer primer and nested CB primers for 55 cycles as described in Example IX.

TCR β -chain sequences of human MBP-reactive T cells

- VB16mer amplified TCR β -chain genes from human MBP-
10 reactive T cell lines were sequenced using the CBseq primer. Amplification products were gel purified, base denatured and sequenced from the CBseq primer. Readable DNA sequence was obtained from 5 of these lines, indicating that predominant T cell clones had been selected by long
15 term in vitro passage. One of these sequences, from the Re cell line (Table VI), possessed a β -chain VDJ amino acid sequence that shared five of the first six and six of nine total residues with the β -chain VDJ amino acid sequence conserved among MBP reactive, encephalogenic T cells in the
20 B10.PL mouse model of EAE. This sequence was not present among the predominant TCR rearrangements found in the remaining four human MBP reactive T cell lines.

- To determine if similar sequences were present in the β -chain repertoire of the MBP-reactive T cell lines from
25 other MS patients, PCR amplification was conducted with a degenerate (n=1024) 21-nucleotide primer (VBRe) corresponding to seven amino acids of this sequence (Table VI). RNAs were reversed transcribed and amplified in 20 cycle stage I reactions with the VB16mer and CBext primers.
30 One μ l aliquots of these stage I reactions were reamplified for 35 cycles with the VBRe and CB int primers. One μ l aliquots of these reactions were analyzed by Southern blot hybridization with a 32 P-labeled human CB probe. This

analysis revealed the 300 bp amplified product in the Re cell line and in one of the other MS patient lines, but not in MBP-reactive T cells from control subjects or in non-MBP reactive human T cell lines and clones. The presence
5 of this sequence in two of the nine MS patient lines tested is compelling. Since this sequence is known to be conserved among encephalogenic T cells in EAE, its detection among MBP-reactive T cells from MS patients demonstrates a role for T cells bearing this determinant in
10 the pathogenesis of MS.

Immunogenic peptides having the sequence SGDQGGNE can be synthesized as shown in Example II and used to immunize human subjects by methods demonstrated in Example III. Such immunizations can result in an effective immune
15 response.

30

A)

TABLE VI

Sample	V β	D β	J β
MS-Re	V β 4.2 ctctgc L C	agcggagaccagggcggc S G D Q G G -	J β 2.1 aatgagcagttcttc N E Q F F
B10.PL		S G D A G G G Y E	

B)

5' G G $\begin{matrix} A \\ C \\ G \\ T \end{matrix}$ G A $\begin{matrix} C \\ T \end{matrix}$ C A $\begin{matrix} A \\ G \end{matrix}$ G G $\begin{matrix} A \\ C \\ G \\ T \end{matrix}$ G G $\begin{matrix} A \\ C \\ G \\ T \end{matrix}$ A A $\begin{matrix} C \\ T \end{matrix}$ G A $\begin{matrix} A \\ G \end{matrix}$ 3'

EXAMPLE IX

Isolation of Oligoclonal Infiltrates of IL-2
Activated VB17 T Cells in the Synovium of
Rheumatoid Arthritis Patients

5 T cell preparations from synovial tissue

Synovial tissue specimens were obtained from radiographically proven rheumatoid arthritis patients undergoing joint replacement therapy. Activated T cells were selected using magnetic beads and antibodies reactive
10 with the human IL2-R (α IL2-R) as follows. Synovial tissue was digested for 4 hrs at 37°C in RPMI + 10% Fetal Bovine Serum (FBS) containing 4 mg/ml collagenase (Worthington Biochemical, Freehold, NJ) and 0.15 mg/ml DNase (Sigma, St. Louis, MO.). Digests were passed through an 80-mesh screen
15 and single cells were collected by Ficoll density gradient centrifugation. Cells at the interface were washed and were incubated at 10^6 /ml for 30 min at 0°C with 5 μ g/ml control mouse IgG (Coulter Immunology, Hialeah, FL) in PBS containing 2% FBS (PBS-FBS). Cells were washed three times
20 and incubated for 30 min at 0°C with magnetic beads conjugated to goat anti-mouse IgG (Advanced Magnetix, Cambridge, MA). Beads were magnetically separated and washed three times with PBS-FBS. This preselection with mouse IgG (mIgG) and magnetic beads was used to control for
25 non-specific adsorption of T cells. The cells remaining in the initial suspension were further incubated 30 minutes at 0°C with 5 μ g/ml monoclonal mouse IgG reactive with the human T cell IL2-R (Coulter Immunology, Hialeah, FL). Cells were washed and selected with magnetic beads as
30 above. Beads from the IgG preadsorption and the IL2-R antibody selection were immediately resuspended in acidified-guanidinium-phenol-chloroform and RNA prepared as described in Chonezynski and Sacchi, Anal. Biochem. 162:156

(1987), which is incorporated herein by reference. Since RNAs were prepared without in vitro culture of the cells and the accompanying bias that may be induced, they are expected to accurately reflect T cell distributions in synovial tissue at the time of surgical removal. Only half of the mIgG and α IL2-R beads from patient 1012 were immediately processed for RNA. The remainder were cultured for 5 days in RPMI 1640, 5% FBS, 20% HL-1 (Ventrex Laboratories Inc., Portland, ME), 25mM HEPES, glutamine, antibiotics and 20% LAK supernatant (Allegretta et al., Science, 247:718 (1990)), which is incorporated by reference herein, as a source of IL-2. RNA was extracted from cultures of the α IL2-R beads (1012IL2.d5), but not from the 1012mIgG sample as no viable cells were present at the end of the 5 day culture.

A T cell clone was derived from the Ficoll pellet of patient 1008. The cells in the pellet were cultured at 2×10^6 /ml in media without IL-2 for two weeks. Non-adherent cells from this culture were cloned by limiting dilution onto autologous synovial cell monolayers. A CD4+ T cell clone 1008.8 was obtained and adapted to culture by regular stimulation with autologous synovial monolayers for 3 days in media without IL-2 followed by a 4 day culture in medium with LAK supernatant.

Lysis of Synovial Adherent Cells by 1008.8

Lysis of synovial adherent cells by 1008.8 was demonstrated as follows. Synovial cell monolayers were labeled as described in Stedman and Campbell, J. Immunol. Meth. 119:291 (1989), which is incorporated herein by reference, with 35 S for use as targets in CTL assays. Cells were trypsinized, washed and plated at 2000 cells per well of a 96-well round bottom microtiter plate. 1008.8 cells, cultured for 3 days prior to the assay with synovial adherent cells and medium containing LAK supernatant, were

added to the targets at the indicated effector:target ratios. Cultures were incubated overnight at 37°C, centrifuged at 300xg for 2 minutes and radioactivity in 50 μ l of the supernatant quantified. Per cent specific lysis was calculated relative to detergent-lysed targets by standard formulas. This clone is cytotoxic for synovial adherent cell targets in CTL assays (Table VII).

TABLE VII

	<u>Effector:Target Ratio</u>	<u>% Specific Lysis</u>
10	5:1	7
	10:1	16
	25:1	32

PCR Amplification of TCR β -chain genes


TCR β -chain genes were amplified with several combinations of the primers shown in Table VIII. The VB16mer primer is a degenerate VB primer (n=256) which is predicted to bind 85% of human TCR β -chain genes at all 16 residues and 95% at 15 residues. This primer has been used to amplify TCR β -chains from more than 25 different human T cell clones, lines or primary tissue preparations. A spectrum of VB genes has been sequenced from these amplified DNAs, arguing against a significant bias of the primer for certain VB families. Thus, PCR amplification with the VB16mer primer facilitates analysis of T cell populations for which a priori knowledge of VB gene usage is unavailable.

T cell receptor β -chain genes were amplified in two-stage amplification reactions with nested pairs of the primers shown in Table VIII. RNAs were reverse transcribed for 1 hour at 42°C with 40pmol of the C β ext primer in a 12 μ l reaction using conditions described by Hart et al., The

Lancet, p. 596 (1988), included by reference herein. Reactions were diluted with a master mix containing 40 pmols of the VB16mer primer, nucleotides and reaction buffer as above but without MgCl_2 to give a final Mg^{+2} concentration of 3.6 mM. Samples were denatured for 15 minutes at 95°C , 1 unit of heat stable recombinant DNA polymerase (Cetus Corporation, Emeryville, CA, Ampli-taqTM) was added and 20 cycles of PCR conducted. Each cycle consisted of a 1 min denaturation at 95°C , a two minute annealing step and a two minute extension at 72°C . The first two cycles were annealed at 37°C and 45°C , respectively, and the remainder at 50°C . One microliter aliquots of these stage I reactions were added to 100 μl stage II amplification reactions (Cetus, Gene-Amp KitTM) containing 100 pmols of the CBint primer and 100 pmols of the VB8, VB17 or 5'CB primers or 700 pmols of the VB16mer primer. Stage II amplifications were conducted as above with a 50°C annealing temperature and without the 37°C and 45°C ramping.

TABLE VIII

A)



B)

V β 16mer	5' T ^G _T T ^A _T C ^C _T T G G T A C ^A _T ^A _C ^A _G ^A _T C A 3'
V β 17	5' T C A C A G A T A G T A A A T G A C T T T C A G 3'
V β 8	5' T C T C C A C T C T G A A G A T C C 3'
5' C β	5' C A A G C T G T T C C C A C C C G A 3'
C β ext	5' C C A G A A G G T G G C C G A G A C 3'
C β int	5' G C G G C T G C T C A G G C A G T A 3'
C β seq	5' C G A C C T C G G G T G G G A A C A 3'

RNA samples from 1012IL2.d5 and 1008.8 cultures were amplified with the VB16mer and CBext primers in stage I reactions and with the VB16mer and the CBint primer in 35 cycle stage II reactions. Reaction products, purified from low melting agarose gel slices with Gene Clean glass beads (Biolol, San Diego, CA), were base denatured and sequenced from the CBseq primer with T7 polymerase (Sequenase, (United States Biochem, Cleveland, OH). A predominate VB sequence, corresponding to a single VB17 rearrangement Table IX, was clearly readable in the 1012IL2.d5 sample. Other, less frequent rearrangements were detected as faint, uninterpretable background bands in the sequencing gels. Culture of these 1012.IL2 beads in IL2-containing medium without added accessory cells or antigen is not expected to induce de novo activation of T cells. Thus, the predominance of a single VB17 rearrangement in this sample reflects in vivo clonal expansion of VB17+ T cells in this patient. DNA sequence determination of TCR β -chain DNA amplified from the cytotoxic T cell clone, 1008.8, also revealed a VB17 rearrangement (Table IX). The presence of VB17 rearrangements in these two different types of synovial T cell samples, derived from two separate RA patients, implicates VB17 bearing T cells in the pathogenesis of RA.

TABLE IX

Sample	V β	D β	J β
1012 day 5	Y L C A S tattctctgtgccagt V β 17	K N P T V S aaaaatcccacgggtctcc	Y G Y T F tatggctacaccttc J β 1.2
1008.8	Y L C A S tattctctgtgccagt V β 17	D N gacaac	E A F F G gaggctttctttgga J β 1.1
1014 IL-2	Y L C A S tattctctgtgccagt V β 17	V R D R R gtgagggacaggaga	N Y G Y T aactatggctacacc J β 1.2
1015 IL-2	Y L C A S S tattctctgtgccagtagt V β 17	S I D S agtatagactcc	S Y E Q Y tcctacgagcagtagt J β 2.7

The presence of VB17 rearrangements in the remaining synovial RNA samples was assessed by PCR amplification with the VB17 specific primer (Table VIII). VB17 TCR DNA was amplified from magnetic bead samples derived from each of the seven RA patients. Ethidium bromide staining of electrophoresed reaction products revealed greater VB17 amplification in four of the α IL2-R samples than in the corresponding mIgG controls. This enrichment was not a product of the isolation procedure, since amplification of VB8 TCRs failed to show differences between mIgG and IL2-R samples.

VB17 rearrangements from two of the α IL2-R RNA preparations were amplified with the VB17 and CBint primer pair and the reaction products sequenced with the CBseq primer. Samples 1014 and 1015 contained single sequences (Table IX), which, like the 1012IL2.d5 sample, demonstrates clonal expansion of VB17 T cells *in vivo*. In contrast, direct sequencing of the rearrangements amplified with the VB8 specific primer was not possible due to significant heterogeneity in the B-chain product.

VB17 has the amino acid sequence MSNQVLCCVVLCLGANTVDG GITQSPKYLFRKEGQNVTLSC EQNLNHDAMYWYRQDPGQGLRLIYYSQIVNDFQKGD IAEGYSVSREKKESFPLTVTSAQKNPTAFYLCASS.


HLA-DR Analysis in Rheumatoid Arthritis Patients

HLA-DR analysis in rheumatoid arthritis patients was performed as follows. DNA from each patient was prepared by boiling 10^5 synovial cells in 200 μ l dH_2O . Ten μ l were amplified for 35 cycles in a 100 μ l reaction (Cetus, Gene Amp KitTM) containing 100 pmols of each of the DRB PCR primers (Table X). One-tenth μ l of this reaction was reamplified in 10 μ ls containing only the DRB2 primer and 17 pmol of α 32P-dCTP as the sole source of dCTP for 10

cycles. Reactions were spiked with 200 μ M dCTP and chased for 2 cycles. The resulting negative strand probes were hybridized to slot blots containing 10 pmol of the HLA-DR allele specific oligos (positive strands) using conditions
5 previously described by Amor et al., J. Immunol. 138:1947 (1987), which is incorporated herein by reference. The slots were washed twice for 20 minutes with tetramethylammoniumchloride (Wood et al., Proc. Natl. Acad. Sci. USA 82:1585 (1985)) which is incorporated herein by
10 reference) at 65-68°C and exposed to X-ray film.

Each of the patients in this study possessed at least one allele of the HLA-DR genes, DR4w4, DR1, DR4w14 or DR4w15, that are known to predispose for RA (Table X).

40



DR β 1 5' G A G T $\overset{A}{\underset{C}{}}$ C T G G A A C A G C 3'

DR β 2 5' G T A G T T G $\overset{A}{\underset{C}{}}$ T T C T G C A 3'

HLA-DR ALLELE-SPECIFIC OLIGONUCLEOTIDES

DRB1 Genes

DR1, DR4w14, DR4w15 5' CTC CTG GAG CAG AGG CGG GCC GCG 3'

DR2 5' T-- --- --- G-C --- ---C --- --- 3'

DR3 5' --- --- --- --- -A- --- -G- CG- 3'

DR4w4 5' --- --- --- --- -A- --- --- 3'

DR4w13 5' --- --- --- --- --- --- --- -A- 3'

DR5, DR6, DR4w10 5' A-- --- --A G-C GA- --- --- --- 3'

DR7 5' A-- --- --- G-C --- --- -G- CA- 3'

DR8 5' T-- --- --A G-C --- --- --- CT- 3'

DR9 5' T-- --- --- -G- --- --- --- -A- 3'

DRB3 Genes

DR2 5' A-- --- --- --- GC- --- --- --- 3'

DR3 5' --- --- --- --- -A- --- -G- CAG 3'

DR7, DR9 5' --- --- --- -G- --- --- --- -A- 3'

B) Patient HLA-DR

1008 1,4w4

1012 1, 3

1013 1, 7

1014 1,4w4

1015 4w4,4w4

1016 N. D.

1017 1, 7

SUBSTITUTE SHEET

T cell receptors containing V β 17 or fragments thereof which are immunogenic or can be made immunogenic can be used to immunize human subjects by methods demonstrated by Example VII. Such immunizations can result in an effective
5 immune response.

Although the invention has been described with reference to the presently-preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention.
10 Accordingly, the invention is limited only by the following claims.

WE CLAIM:

1. A vaccine for preventing or treating a T cell mediated pathology or an unregulated T cell clonal replication in a mammal, comprising an immunogenically effective amount of a T cell receptor or fragment thereof corresponding to a T cell receptor present on the surface of T cells mediating said pathology, and a pharmaceutically acceptable medium.

2. The vaccine of claim 1, wherein said T cell mediated pathology is rheumatoid arthritis and wherein said T cell receptor comprises the amino acid sequence of VB17.

3. The vaccine of claim 1, wherein said fragment comprises a variable region sequence of said T cell receptor.

4. The vaccine of claim 3, wherein said variable region sequence is the β -chain variable region.

5. The vaccine of claim 4, wherein said T cell mediated pathology is rheumatoid arthritis and wherein said β -chain variable region comprises substantially the amino acid sequence designated VB17.

6. The vaccine of claim 5, wherein said β -chain variable region comprises substantially the sequence SQIVNDFQK.

7. The vaccine of claim 1, wherein said fragment comprises a V(D)J junctional sequence.

8. The vaccine of claim 1, wherein said fragment comprises a junctional region sequence.

9. The vaccine of claim 1, further comprising an adjuvant.

10. The vaccine of claim 1, wherein said vaccine comprises more than one type of T cell receptor or fragment thereof.

11. The vaccine of claim 1, wherein said vaccine comprises more than one fragment corresponding to different sequences of the same T cell receptor.

12. The vaccine of claim 1, wherein said fragment is conjugated to a carrier.

13. The vaccine of claim 1, wherein said T cell mediated pathology is multiple sclerosis, said mammal is a human, and said fragment comprises substantially the sequence SGDQGGNE so as to elicit an immune response against T cell receptors having substantially the sequence SGDQGGNE.

14. The vaccine of claim 1, wherein said T cell receptor comprises the sequence SGDQGGNE.

15. A method of vaccinating an individual exhibiting or at risk of exhibiting a T cell-mediated pathology, comprising administering to the individual the vaccine of claim 1.

16. The method of claim 15, wherein said vaccine is administered more than once.

17. The method of claim 15, wherein said vaccine is administered in a formulation including an adjuvant.

18. A method of treating unregulated T cell clonal replication in an individual, comprising administering the vaccine of claim 1 to the individual.

19. A method of selecting a vaccine for use in treating a T cell mediated pathology comprising the steps of:

a. obtaining T cell clones mediating said condition;

b. determining the amino acid sequence of T cell receptors from T cell clones associated with said condition;

c. selecting segments of those T cell receptors which are characteristic of said associated T cell receptors but not of non-associated T cell receptors; and

d. selecting amino acid sequences of said selected sequences which are capable of eliciting an immunogenic response to said T cell receptor, and thereby selecting the vaccine.

20. A method of diagnosing or predicting susceptibility to rheumatoid arthritis in an individual comprising detecting T cells having the β -chain variable region designated VB17 or a fragment thereof in a sample from the individual, the presence of abnormal expression of VB17-containing T cells indicating rheumatoid arthritis or susceptibility to rheumatoid arthritis.

21. The method of claim 20, comprising detecting a portion of said VB17 which substantially does not occur on non-rheumatoid arthritis associated T-cell receptors.

22. The method of claim 20, wherein said sample is from synovial tissue.

23. The method of claim 20, wherein said VB17 is detected by contacting said VB17 with a detectable ligand.

24. The method of claim 20, wherein the presence of said VB17 is detected by a nucleotide probe which is complementary to the nucleotide sequence encoding VB17.

25. A method of preventing or treating rheumatoid arthritis comprising preventing the attachment of a VB17 containing T-cell receptor to its binding partner.

26. The method of claim 25, wherein said binding partner is an HLA-DR predisposing for rheumatoid arthritis.

27. The method of claim 25, wherein attachment is prevented by binding a ligand to VB17.

28. The method of claim 25, wherein attachment is prevented by binding a ligand to said VB17 binding partner.

29. A method of preventing or treating rheumatoid arthritis in an individual comprising cytotoxically or cytostatically treating VB17 containing T-cells in the individual.

30. The method of claim 29, wherein said VB17 containing T-cells are treated with a cytotoxic or cytostatic agent which selectively binds VB17.

31. The method of claim 30, wherein said agent is an antibody attached to a moiety selected from the group consisting of radioactive moieties, chemotherapeutic moieties and chemotoxic moieties.

32. A vaccine for preventing or treating a T cell mediated pathology or unregulated T cell clonal replication in a mammal, comprising anti-idiotypic antibodies which are internal images of a T cell receptor or a fragment thereof corresponding to a cell receptor present on the surface of a T cell mediating said pathology and a pharmaceutically acceptable medium.

33. The vaccine of claim 32, wherein said fragment comprises a variable region sequence of the T cell receptor.

34. The vaccine of claim 32, wherein said fragment comprises a V (D) J junction sequence.

35. The vaccine of claim 32, further comprising an adjuvant.

36. The vaccine of claim 32, wherein said vaccine comprises anti-idiotypic antibodies which are internal images of more than one T cell receptor or fragment thereof.

37. A method of vaccinating an individual exhibiting or at risk of exhibiting a T cell-mediated pathology or an unregulated T cell clonal replication, comprising administering to the individual the vaccine of claim 32.

38. The vaccine of claim 32, wherein said fragment comprises a junctional region sequence.

39. A method of diagnosing or predicting susceptibility to multiple sclerosis in an individual comprising detecting T cells having substantially the sequence SGDQGGNE in a sample from the individual, the presence of such T cells indicating multiple sclerosis or susceptibility to multiple sclerosis.

40. The method of claim 39, wherein said sequence is detected by contacting said sequence with a detectable ligand.

41. The method of claim 39, wherein the presence of said sequence is detected by a nucleotide probe which is complementary to the nucleotide sequence encoding said sequence.

42. A method of preventing or treating multiple sclerosis comprising preventing the attachment of a T-cell receptor containing substantially the sequence SGDQGGNE to its binding partner.

43. The method of claim 42, wherein attachment is prevented by binding a ligand to said sequence.

44. The method of claim 42, wherein attachment is prevented by binding a ligand to said T cell receptor binding partner.

45. A method of preventing or treating multiple sclerosis in an individual comprising cytotoxicly or cytostatically treating T cells containing substantially the sequence SGDQGGNE in the individual.

46. The method of claim 45, wherein said sequence containing T cells are treated with a cytotoxic or cytostatic agent which selectively binds said sequence.

47. The method of claim 45, wherein said agent is an antibody attached to a moiety selected from the group consisting of a radioactive moieties and a chemotherapeutic moieties and chemotactic moieties.


48.

48. A composition of matter comprising a T cell receptor or a fragment thereof corresponding to a T cell receptor present on the surface of T cells mediating a pathology and an adjuvant.

49. A peptide comprising the sequence SGDQGGNE.

INTERNATIONAL SEARCH REPORT

International Application No PCT/US 90/01516

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ : C 07 K 7/06 , A 61 K 37/02, G 01 N 33/564, G 01 N 33/68		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁵	C 12 N, C 07 K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with Indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	WO, A, 86/06413 (CALIFORNIA INSTITUTE OF TECHNOLOGY) 6 November 1986 see the whole document --	1,7-12,19, 32,35-36,38, 48
A	Nature, volume 308, 8 March 1984, Y. Yanagi et al.: "A human T cell-specific cDNA clone encodes a protein having extensive homology to immunoglobulin chains", see pages 145-149 --	
A	Research Immunology, volume 140, no. 2, 1989, W.E. Biddison et al.: "The germline repertoire of T-cell receptor beta-chain genes in patients with multiple sclerosis", see pages 212-215 -- ./.	
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
5th July 1990		17. 08. 90
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		 M. SOTELO

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, " with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	Immunol. Res., volume 8, no. 2, 1989, C. Richard Ross et al.: "Antibodies to synthetic peptides corresponding to variable-region first-framework segments of T cell receptors", see pages 81-97 --	
A	Chemical Abstracts, volume 105, no. 1, 7 July 1986, (Columbus, Ohio, US), S.F. Schluter et al.: "Antibodies to synthetic joining segment peptide of the T-cell receptor β -chain: serological cross-reaction between products of T-cell receptor genes, antigen binding T-cell receptors, and immunoglobulins", see page 464, abstract 4767q, & Proc. Natl. Acad. Sci. USA 1986, 83(6), 1872-6 (Eng). --	
X,P	Science, volume 246, no. 4930, 1989, M.D. Howell et al.: "Vaccination against experimental allergic encephalomyelitis with T cell receptor peptides", pages 668-670 see page 668-670; tables 1 & 2 --	1,9,19,32,38, 48
X,P	Letters to Nature, volume 341, no. 6242, 12 October 1989, A.A. Vandembark et al.: "Immunization with a synthetic T-cell receptor V- region peptide protects against experimental autoimmune encephalomyelitis", pages 541-544 see the whole document -----	1,3,4,9,11,12, 19,32-33,35- 36,48

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers because they relate to subject matter not required to be searched by this Authority, namely:
 **claims searched completely: 1-14, 19-24, 32-36, 38-41,
 48, 49
 claims not searched: 15-18, 25-31, 37, 42-47

see PCT rule 39.1(iv): method for treatment of the human
or animal body by therapy

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers....., because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING:

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.

US 9001516
SA 35869

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

E.O. FORM 10479